

2011 Fall Avian Survey

Wilton IV Wind Energy Center
Burleigh County, North Dakota



Prepared for



June 2012



TETRA TECH EC, INC.

EXECUTIVE SUMMARY

Tetra Tech EC, Inc. (Tetra Tech) was contracted by Wilton Wind IV, LLC (Wilton IV), a subsidiary of NextEra Energy Resources, LLC (NextEra), to undertake fall avian use surveys for the proposed Wilton IV Wind Energy Center (Project) in Burleigh County, North Dakota. The studies were conducted to identify potential avian impacts associated with building and operating the Project. Birds have been identified as a group potentially at risk because of collisions with wind turbines and power lines and displacement due to the presence of the associated structures. Weekly surveys were performed for the Project from August 9 to November 11, 2011, which included the fall through early winter seasons. Fixed point count surveys (800-meter [m] radius) were conducted at 7 points distributed throughout the Project Area, which encompasses 15,752 acres of privately owned predominantly agricultural land.

A total 4,083 birds were observed within the Project Area, comprised of 3,087 birds from 42 species and 996 birds that could not be identified to species. Overall mean bird use within the Project Area was 41.66 birds/20 minutes (min) with individual 20-min surveys ranging 0 to 600 birds/20 min.

Four non-raptor species and one unidentified species were the most frequently observed within the Project Area. The non-raptors with the highest mean use were unidentified blackbird (may include red-winged blackbird, common grackle, and brown-headed cowbird; 10.15 birds/20 min), red-winged blackbird (8.47 birds/20 min), horned lark (8.23 birds/20 min), snow goose (5.51 birds/20 min), and common grackle (2.74 birds/20 min). The unidentified blackbirds had the second highest encounter rate of 5.05 birds flying at RSA height/20 min. The red-winged blackbird, common grackle, snow goose, and horned lark are all widespread species. These species also had high encounter rates (7.16, 1.19, 1.17, and 0.82 birds flying at RSA height/20 min, respectively) within the anticipated RSA but also have large regional breeding populations in North Dakota (red-winged blackbird, common grackle, and horned lark) and northern Canada (snow geese). If fatalities do occur they are unlikely to have population-level consequences.

Raptors are a group of special interest because of their propensity to fly at heights similar to a turbine RSA. Overall mean use for raptors was 0.35 birds/20 min. Red-tailed hawks and northern harriers were the most common raptors observed in the Project Area, but in low numbers (0.17 and 0.10 birds/20 min). Red-tailed hawk had a low encounter rate (0.08 birds flying at RSA height/20 min) and northern harrier had no encounter rate within the intended RSA. As a result any potential for turbine-related impacts to these species is anticipated to be low. Additional raptor species observed during point counts included: Swainson's hawk, American kestrel, unidentified buteo, sharp-shinned hawk, and broad-winged hawk.

LISTED AND SENSITIVE SPECIES

No federally listed avian species were observed during fall avian surveys or as an incidental observation within the Project Area.

At the state level, North Dakota does not have a list defined by statute for threatened or endangered species comparable to the federal Endangered Species Act (ESA). North Dakota has instead identified 100 Species of Conservation Priority under the State Wildlife Action Plan.

State-designated Species of Conservation Priority observed in the Project Area include grasshopper sparrow (level I), Swainson's hawk (level I), ferruginous hawk (level I), northern harrier (level II), and sharp-tailed grouse (level II). The designation of Species of Conservation Priority describes a species identified as in decline at the national, regional or state level, or a species whose population status is not well known, but thought to be in decline in North Dakota. Species of Conservation Priority receive special attention from state agencies, but do not require take permits or have other regulatory implications regardless of status (Level I or II), but all of these species are protected under the Migratory Bird Treaty except sharp-tailed grouse. The sharp-tailed grouse is managed as a game-bird in North Dakota. All species listed above had no encounter rates within the intended RSA, primarily because of their low occurrence within the Project Area.

Table ES-1. Fall avian use summary

| Variable | Result | Details |
|---|--------------------|--|
| Non-raptors | | |
| Mean use | 41.32 birds/20 min | |
| Number of species with high encounter rates (>1.0 birds at RSA height/20 min) | 3 | red-winged blackbird, common grackle and snow goose (Section 3.2) |
| Federally listed ¹ species observed within the Project Area | No | |
| State-listed species ² within the Project Area | Yes | grasshopper sparrow and sharp-tailed grouse (Section 4.3) |
| State-listed species within RSA | No | |
| Raptors | | |
| Mean use | 0.35 birds/20 min | |
| Number of species with high encounter rates (>1.0 birds at RSA height/20 min) | None | |
| Eagles observed within the Project Area | No | |
| Federally listed species observed within the Project Area | No | |
| State-listed species within the Project Area | Yes | Swainson's hawk, ferruginous hawk and northern harrier (Section 4.3) |
| State-listed species within the RSA | No | |
| Habitat | | |
| Native habitat likely to be affected by development | Yes | Native prairie |
| Lakes (waterfowl attractant) | Yes | Small cattle ponds |
| Wetlands (attractant for cranes, waterfowl, and other water-based species) | Yes | Scattered throughout |
| Cliffs (raptor nesting and traveling) | No | |
| River (permanent water source, migration corridor) | Yes | West branch of Apple Creek |
| Known refuges or habitat features that may funnel migrants | None | |

¹Federally listed species include species listed as endangered, threatened, or candidate species in the Endangered Species Act.

²The North Dakota Game and Fish Department maintains a list of Species of Conservation Priority (Hagen et al. 2005) but there are no permitting requirements for listed species. State species listed are those in addition to federally listed species.

TABLE OF CONTENTS

| | |
|--|-------------|
| EXECUTIVE SUMMARY | ES-1 |
| 1.0 INTRODUCTION | 1 |
| 1.1 WIND ENERGY AND BIRDS | 1 |
| 1.2 STUDY DESCRIPTION | 2 |
| 2.0 METHODS | 2 |
| 2.1 AVIAN SURVEYS | 3 |
| 2.1.1 Fixed-point Surveys | 3 |
| 2.1.2 Incidental Observations | 4 |
| 2.1.3 Listed Species Information | 4 |
| 2.1.4 Data Quality Assurance/Quality Control | 4 |
| 2.2 ANALYSIS | 5 |
| 2.2.1 Species Groupings | 5 |
| 2.2.2 Avian Use of the Wilton IV Project | 5 |
| 2.2.3 Flight Behavior | 5 |
| 2.2.4 Encounter Rate | 5 |
| 2.2.5 Mortality Risk | 6 |
| 3.0 RESULTS | 6 |
| 3.1 AVIAN USE AND FREQUENCY OF OCCURRENCE. | 6 |
| 3.2 FLIGHT HEIGHT AND ENCOUNTER RATE | 8 |
| 3.3 LISTED SPECIES | 8 |
| 3.6 INCIDENTAL OBSERVATIONS | 8 |
| 4.0 DISCUSSION AND RECOMMENDATIONS | 9 |
| 4.1 NON-RAPTOR USE AND ENCOUNTER RATE | 9 |
| 4.2 RAPTOR USE AND ENCOUNTER RATE | 10 |
| 4.3 LISTED AND SENSITIVE SPECIES | 10 |
| 4.4 WILTON IV PROJECT CONCLUSIONS | 10 |
| 5.0 REFERENCES | 12 |

LIST OF FIGURES

- Figure 1. Wilton IV Wind Energy Center Vicinity map.
Figure 2. Wilton IV Wind Energy Center Point count location map (Fall 2011).
Figure 3. Non-raptor mean use by survey date in Fall 2011 at the Wilton IV Wind Energy Center.
Figure 4. Non-raptor mean use by point count location (Fall 2011).
Figure 5. Raptor mean use by survey date in Fall 2011 at the Wilton IV Wind Energy Center.
Figure 6. Raptor mean use by point count location (Fall 2011).

LIST OF TABLES

- Table ES-1. Fall avian use summary.
Table 1. Fall 2011 point count survey dates at the Wilton IV Wind Energy Center..
Table 2. Avian species, by species grouping, observed during Fall 2011 point count surveys at the Wilton IV Wind Energy Center.
Table 3. Avian species observed by point during Fall 2011 point count surveys at the Wilton IV Wind Energy Center.
Table 4. Summary of avian flight heights in relation to the turbine rotor swept area (RSA) during Fall 2011 point count surveys at the Wilton IV Wind Energy Center.
Table 5. Avian flight height characteristics in relation to the turbine rotor swept area (RSA) during Fall 2011 point count surveys at the Wilton IV Wind Energy Center.
Table 6. Incidental observations of birds during Fall 2011 point counts at the Wilton IV Wind Energy Center.

LIST OF APPENDICES

- Appendix 1. Flight directions of birds observed during Fall 2011 point count surveys at the Wilton IV Wind Energy Center.

1.0 INTRODUCTION

1.1 WIND ENERGY AND BIRDS

Wind energy provides a clean, renewable energy source that is in high demand. As wind power has become more common, the need to address potential environmental impacts has increased. Birds have been identified as a group potentially at risk because of collisions with wind turbines and power lines and displacement due to the presence of the associated structures (Erickson et al. 2005, Drewitt and Langston 2006, Arnett et al. 2007). Specifically, migrant passerines (e.g., songbirds) are found more often in post-construction mortality monitoring compared to other groups of birds (Arnett et al. 2007). In fact, at newer generation wind energy facilities outside of California, approximately 80 percent of documented mortalities have been songbirds, of which 50 percent are often nocturnal migrants (Erickson et al. 2001, Drewitt and Langston 2006, Johnson et al. 2007, Strickland and Morrison 2008). Data based on radar data and mortality monitoring suggest that less than 0.01 percent of migrant songbirds that pass over wind farms are killed (Erickson 2007). Locally breeding songbirds may experience lower mortality rates than migrants because many of these species tend not to fly at turbine heights during the breeding season. However, some breeding songbird species have behaviors that increase the risk of collisions with turbines. For example, horned larks have been commonly found as fatalities at wind farms (Erickson et al. 2002). Mortality may be partially attributed to the territorial flight displays in which male horned lark fly to heights of 80 meters (m) to 250 m (Pickwell 1931).

Despite the observation that most wind farm fatalities are songbirds, raptor mortality historically has received the most attention. Raptor mortality at newer generation wind projects has been low relative to previous generation wind farms, although there is substantial regional variation (Erickson et al. 2002, 2004, Johnson et al. 2002, Kerns and Kerlinger 2004, Jain et al. 2007). Although raptor mortality is reduced at newer generation facilities, mortality may not be eliminated by advances in turbine technology (e.g., turbine height, tower structure) and local micro-siting and site evaluation efforts are still necessary.

In addition to mortality associated with wind farms, concerns have been raised that some bird species may avoid areas near turbines after a wind farm is in operation (Drewitt and Langston 2006). For example, at the Buffalo Ridge wind energy facility in Minnesota, densities of male songbirds were significantly lower in Conservation Reserve Program (CRP) grasslands containing turbines than in CRP grasslands without turbines. It was suggested that the reduced density may be due to avoidance of turbine noise and maintenance activities, and reduced habitat quality due to the presence of access roads and large gravel pads surrounding the turbines (Leddy et al. 1999). Reduced abundance of grassland songbirds was found within 50 m of a turbine pad for a wind farm in Washington and Oregon, but the investigators attributed displacement to the direct loss of habitat or reduced habitat quality and not the presence of the turbines (Erickson et al. 2004). Research at two sites in North and South Dakota (Shaffer and Johnson 2008) suggests that certain grassland songbird species (two of four studied) may avoid turbines by as much as 200 m but these results have not been finalized nor verified at additional sites. None of these studies have addressed whether or not these avoidance effects are temporary (i.e. the birds may habituate to the presence of turbines over time) or permanent.

Finally, most native, migratory birds are protected under the Migratory Bird Treaty Act (MBTA) of 1918. Under the MBTA it is unlawful to pursue, hunt, take, capture or kill; attempt to take, capture or kill; possess, offer to or sell, barter, purchase, deliver or cause to be shipped, exported, imported, transported, carried or received any migratory bird, part, nest, egg or product. Despite extensive liability provisions, the USFWS has narrowly interpreted its permitting authority. “As currently written, USFWS’s regulations establish a permitting scheme for a variety of intentional activities, such as hunting, falconry, certain import and export activities, depredation control, and scientific research. But...there is no permitting scheme for the incidental take of migratory birds during otherwise lawful activities” (Beveridge 2005). There is no permitting framework (i.e., incidental take permits) that allow a wind company to protect itself from liability at wind facilities; however, the USFWS does not usually take action if good faith efforts have been made to minimize impacts. To date, no wind development company has been charged for violations of the MBTA.

1.2 STUDY DESCRIPTION

Wilton Wind IV, LLC (Wilton IV), a subsidiary of NextEra Energy Resources, LLC, is planning to develop the Wilton IV Wind Energy Center (Project) in Burleigh County, North Dakota (Figure 1). Wilton IV is committed to environmental due diligence and has contracted Tetra Tech EC, Inc. (Tetra Tech) to conduct fall avian surveys in the Project Area to quantify local avian use in the area and to identify potential avian impacts associated with building and/or operating the proposed facility.

The Project Area encompasses 15,752 acres and is located in the Northwestern Glaciated Plains Ecoregion, with western portions of the Project Area located in the Northwestern Great Plains Ecoregion (Bryce et al. 1996). Historically, much of the landscape was a mix of western mixed-grass prairie, short-grass prairie, with associated wetlands of the Missouri Slope and River Breaks subregions (Bryce et al. 1996). This semiarid region of North Dakota includes level to rolling plains topography with isolated sandstone buttes or badlands formations. Today, native grasslands still persist in mostly areas of steep or broken topography and have been largely been replaced by agriculture. Land use is predominantly dry-land farming of barley, sunflowers, corn, spring and winter wheat interspersed with cattle grazing.

North Dakota has over 365 documented bird species (Faanes and Stewart 1982) and is situated within the Central Flyway, one of the main bird migratory routes (USFWS 2011a). The Central Flyway runs through the central portion of the U.S. and, as a consequence, the Project Area. Most birds that move along the Central Flyway travel from Canada through the central states, eventually reaching the tropics of South America via the Gulf of Mexico (USFWS 2011a).

2.0 METHODS

To evaluate avian risk at wind energy facilities, standardized protocols for pre-construction point counts have been established and were used in this study. This protocol is designed to be responsive to the level of effort recommended in the National Wind Coordinating Committee’s *Comprehensive Guide to Studying Wind Energy/Wildlife Interactions* (Strickland et al. 2011) and the *U.S. Fish and Wildlife Service Draft Land-Based Wind Energy Guidelines* (USFWS 2011b). Data collected from these counts are used to identify species or species groups that may be at

risk from project development and may provide additional information for micro-siting wind facilities to minimize impacts to birds. Results in this report are presented in terms of species groups, and highlight federally and state-listed species, and species of concern.

2.1 AVIAN SURVEYS

2.1.1 Fixed-point Surveys

Experienced field biologists conducted 20-minute (min) point count surveys at 7 locations within the Project Area to evaluate avian use, behavior, and species composition during fall migration (Figure 2). The biologists conducted weekly surveys from August 9 to November 11 (Table 1), thereby encompassing the fall migration to early winter seasons. Tetra Tech distributed the survey locations throughout the Project Area and chose locations that maximized the 360-degree sight distance for the observer and covered a diversity of habitats.

The field biologists collected data on all birds observed within an 800-m radius of the point count location. Surveys at each point lasted for 20 min, during which time biologists continuously recorded any visual or auditory observations. Biologists recorded the following data: species, number of individuals, time of observation, height aboveground, behavior, and flight direction. Flight direction data was only recorded for those birds that were exhibiting clear migratory behavior, and not for those birds that were making localized flights within the Project Area. Data on flight direction can be found in Appendix 1. The biologists estimated flight heights and distances using existing meteorological towers, local transmission lines, and topographic maps for reference.

The survey protocol used in this study is designed to collect data on all bird species and to provide results that are comparable with other studies at wind farms rather than to target specific taxa. The benefit of using this protocol is that it estimates avian use throughout the day and captures activity by a variety of bird species. During the breeding season, and to a lesser extent in the fall and winter, songbirds are most active in the morning and can be difficult to detect during the afternoon. In contrast, raptors become active as the sunlight heats the air and creates thermals, which individuals use for soaring (Ballam 1984). Thus, raptors are more readily detected several hours after sunrise. Therefore, this protocol is appropriate for characterizing the bird community using the Project Area during this time of year.

Tetra Tech chose 20-min survey periods because they provide adequate time to detect both raptors and non-raptors. However, time periods of 20 min may lead to double-counting of songbirds (i.e., counting the same individual more than once) because individuals may appear and disappear from view. For example, if a horned lark is detected perched on a fence then disappears from view and, 6 minutes later, a horned lark is seen flying, these birds are recorded as separate observations because it is not possible to distinguish individuals. Double-counting of birds is not problematic for this type of survey because the objective is to document use in terms of number of birds noted per 20-min survey, not number of distinct individual birds.

Detectability varies among species and potentially not all individuals within the 800-m radius were counted. This variation in detectability results in an overestimate of mean use for conspicuous species and an underestimate of mean use for reclusive species (Thompson 2002).

Birds not easily identifiable, such as those seen under low light conditions or small birds seen at a distance were identified to the lowest taxonomic level possible. Hence, unidentified birds are included in the results.

2.1.2 Incidental Observations

Incidental observations included observations that occurred 1) during travel between points, 2) before or after the official 20-min survey period, and 3) outside of the 800-m radius circular plot. Biologists recorded these observations on separate data sheets and these data were not used in the formal analysis; however, a summary of incidental birds is presented to provide additional information about species found in the local area.

2.1.3 Listed Species Information

A list of species currently protected under the Endangered Species Act (ESA) can be found at <http://www.fws.gov/angered/>. Under the ESA, it is unlawful for a person to take a listed animal without a permit. Take is defined as “to harass, harm, pursue, hunt, shoot, wound, kill, trap, capture, or collect or attempt to engage in any such conduct.” Through regulations, the term “harm” is defined as “an act which actually kills or injures wildlife. Such an act may include significant habitat modification or degradation where it actually kills or injures wildlife by significantly impairing essential behavioral patterns, including breeding, feeding, or sheltering.”

The North Dakota Game and Fish Department (NDGFD) have identified 100 Species of Conservation Priority within North Dakota. These species are ranked in three priority levels based on such factors as known status, funding availability, and presence of breeding habitat within North Dakota (Hagen et al. 2005). The definitions of each rank are listed below:

- Level I: A species having a high level of conservation priority because of declining status either in North Dakota or across their range; or a high rate of occurrence in North Dakota constituting the core of the species’ breeding range, but are at-risk range wide, and non-State Wildlife Grants funding is not readily available to them.
- Level II: Species having a moderate level of conservation priority; or a high level of conservation priority, but a substantial amount of non-State Wildlife Grant funding is available to them.
- Level III: North Dakota’s species having a moderate level of conservation priority, but are believed to be peripheral or do not breed in North Dakota.

Species that are listed under the 100 Species of Conservation Priority are not afforded any formal protection by the state or require special take permits. Additional information on North Dakota Species of Conservation Priority can be found at: <http://www.gf.nd.gov/conservation/levels-list.html>.

2.1.4 Data Quality Assurance/Quality Control

Tetra Tech implemented quality assurance and quality control measures during all stages of data collection, analysis, and report preparation. To ensure legibility and completeness of data sheets,

each biologist reviewed, and clarified if needed, all data sheets before data entry into a FileMaker Pro™ relational database for data storage and analysis. Prior to analysis, an independent reviewer conducted a 100-percent quality review of the data entries. Any questions that arose at this time were directed toward and answered by the field biologists.

2.2 ANALYSIS

2.2.1 Species Groupings

Tetra Tech considered two primary groups of interest: raptors and non-raptors. Tetra Tech defined raptors as vultures, hawks, eagles, falcons, and owls. As turkey vulture flight behavior is similar to raptors and as they are often included as raptors in other studies, Tetra Tech has included them with raptors for the purpose of analysis. Non-raptors were defined as all other species groups.

2.2.2 Avian Use of the Wilton IV Project Area

Tetra Tech derived avian use (mean use) of the Project Area by calculating the average number of birds observed per 20-min survey at each point. To evaluate the diversity and composition of avian species using the Project, Tetra Tech first summarized the number of individuals (birds/20 min) and species. Tetra Tech also calculated a measure of variability (90 percent confidence intervals) for all mean use values. In addition, the number of observations is also presented, where an observation can be either an individual bird or a discrete flock of birds. This information helps evaluate whether high mean use is driven by a single event (e.g., a large flock of birds moving through the Project Area on migration). Because individual birds are not uniquely marked and identified, actual population size or abundance cannot be determined. One individual may be counted multiple times during a survey period or across survey periods. Therefore, avian mean use does not equate to abundance.

2.2.3 Flight Behavior

Tetra Tech evaluated flight behavior by calculating the proportion of flying birds observed below, within, or above the height of the anticipated turbine rotor swept area (RSA). Wilton IV proposes to use the GE 1.6 MW-100 wind turbine for the Project; therefore, a turbine size with an 80-m hub height and 100-m rotor diameter was used to calculate the RSA. With these specifications, the estimated RSA was between 30.0 and 130.0 m above ground. Tetra Tech considered a bird to have flown within the height of the anticipated RSA if any of its recorded heights fell within the upper or lower limits of the anticipated RSA.

2.2.4 Encounter Rate

To estimate the rate at which a species flew at the height of the anticipated RSA, Tetra Tech applied the following equation to every species observed in the Project Area:

$$\text{Encounter Rate} = A * P_f * P_t$$

A is the mean number of birds/20 min for a given species, P_f is the proportion of all activity observations for a given species that were flying; and P_t is the proportion flying observations that were at the height of a turbine RSA for a given species. The encounter rate provides information

on the rate at which a species may move at a height that is consistent with the RSA of the proposed turbines. This information is an important component in evaluating risk of collisions; however, this number alone does not indicate risk to a species. Species with a high encounter rate are at a higher risk of collision than species with a low encounter rate, but it does not mean that mortality is certain. Other factors such as turbine location or a species ability to detect turbine blades, flight maneuverability, and habitat selection also influence mortality (Orloff and Flannery 1992). Values are sensitive to large flocks of birds flying within the RSA; that is, a species will have a high encounter rate even if only seen a few times in large flying flocks. Encounter rate also does not account for migrating behavior of nocturnal migrants.

2.2.5 Mortality Risk

The relationship between pre-construction avian use and post-construction mortality is not yet completely defined due to a lack of pre- and post-construction data from sites with moderate to high use, although a recent meta-analysis suggests that pre-construction studies provide poor indicators of post-construction mortality (Ferrer et al. 2011). Based on the available data, raptor fatality rates generally are low at most wind energy developments with exceptions demonstrated at certain facilities in California with a predominance of older generation turbines (NWCC 2010). The highly regional nature of avian mean use across North America and the scarce data on avian mortality at wind farms in many parts of the continent contribute to uncertainty in predicting fatality rates (Arnett et al. 2007). To date, the most comprehensive source of regional information on avian fatality rates is the *Avian and Bat Fact Sheet* which shows that estimated fatalities range from 14 birds/MW/year in Tennessee to 0 birds/MW/year in Oklahoma (NWCC 2010). As a result of uncertainty in predicting fatality rates, Tetra Tech did not attempt to derive mortality estimates from mean use data but will highlight those species or groups that may experience mortality or displacement that could significantly affect local or regional populations, based on the data provided in this report and other information sources.

3.0 RESULTS

3.1 AVIAN USE AND FREQUENCY OF OCCURRENCE.

Biologists surveyed 3,476 acres of the Project Area during point count surveys, covering 22 percent of the total Project Area. The 7 point count locations were surveyed 14 times, resulting in 98 total 20-min surveys. A total 4,083 birds were observed within the Project Area, comprised of 3,087 birds from 42 species and 996 birds that could not be identified to species during the 98 fixed-point count surveys (Table 2). Overall mean bird use within the Project Area was 41.66 birds/20 min and ranged from 0 to 600 birds/20 min survey.

Overall mean use by non-raptors was 41.32 birds/20 min and, among species groups, mean use was highest for songbirds (33.80 birds/20 min; Table 2). Songbirds were observed in the majority of surveys and were widely distributed throughout the Project Area. The songbirds with the highest mean use were unidentified blackbird (10.15 birds/20 min, observed in 10.2 percent of all surveys) red-winged blackbird (8.47 birds/20 min; 14.3 percent of all surveys) and horned lark (8.23 birds/20 min; 40.8 percent of all surveys – Table 2). These two species and the unidentified blackbird comprised 79.5 percent of the songbird species group and 64.5 percent of all birds observed (Table 2). Additional songbird species with a high frequency included American goldfinch (18.4 percent), barn swallow (16.3 percent), common grackle (14.3 percent),

and western meadowlark (11.2 percent). Every other songbird species was detected in less than 10 percent of all surveys.

Among waterfowl, the second highest mean use for species groups (5.91 birds/20 min), the species with a relatively high mean use value was the snow goose (5.51 birds/20 min, was observed in 3.1 percent of all surveys in 3 flocks; Table 2). Snow geese accounted for 93.3 percent of the waterfowl species group.

The remaining species groups, pigeons/doves, raptors, gamebirds, waterbirds, cranes/rails, and woodpeckers had the next highest mean use values in order (0.87, 0.35, 0.33, 0.27, 0.12, and 0.03 birds/20 min respectively; Table 2).

Non-raptor mean use was highest on October 28 (184.29 birds/20 min) and October 20 (103.71 birds/20 min; Figure 3). The species that contributed to high mean use on October 28 were the snow goose (540 individuals in 3 flocks - largest flock was 250 individuals) and unidentified blackbirds (250 individuals all in one flock). The species that contributed to high mean use on October 20 was the red-winged blackbird (665 individuals in 5 flocks – largest flock was 500 individuals). Mean use for non-raptors was highest at point 4 (80.14 birds/20 min) and observations at this point included red-winged blackbirds (633 individuals) and unidentified blackbirds (298 individuals; Table 3; Figure 4). Mean use for non-raptors was also high for point 2 and point 7 (71.71 and 70.21 birds/20min respectively; Figure 4). Observations at point 2 included unidentified blackbird (265 individuals), horned lark (232 individuals), common grackle (118 individuals), and red-winged blackbird (111 individuals, Table 3). Observations at point 7 included unidentified blackbirds (350) horned larks (287) and snow geese (250, Table 3). The snow geese were observed in flight passing over the Project Area without landing in the immediate area. Habitat at point 4 contained several small wetlands and a large field of unharvested sunflowers that may attract a greater number of blackbirds within the Project Area. Habitat at points 2 and 7 was mostly agriculture fields of wheat and corn and are not considered unique to the Project Area that would concentrate non-raptors to those specific areas.

Raptors are a group of special interest because of their propensity to fly at heights similar to a turbine RSA. Overall mean use for raptors was 0.35 birds/20 min (Table 2) and raptors were the third most frequently observed species groups during the fall surveys. The raptors with the highest use were the red-tailed hawk (0.17 birds/20 min; observed on 14.3 percent of surveys; 0.4 percent of all birds observed) and the northern harrier (0.10 birds/20 min; observed during 10.2 percent of surveys and comprising 0.2 percent of all observations; Table 2). Mean use for each other raptor species was 0.02 birds/20 min or fewer: Swainson's hawk, American kestrel, unidentified buteo, sharp-shinned hawk, and broad-winged hawk.

Mean use by raptors was highest on September 30 (1.14 birds/20 min) and included 4 red-tailed hawks, 2 northern harriers, 1 broad-winged hawk, and 1 sharp-shinned hawk (Figure 5, Table 3). Mean use by raptors was also high for October 15 (1.00 birds/20 min) and included 4 red-tailed hawks, 1 northern harrier, and 1 Swainson's hawk (Figure 5 and Table 3). Mean use by raptors was lower than 0.44 birds/20 min for all other survey dates. Mean use by raptors was highest at point count locations 4 and 2 (0.57 and 0.43 birds/20 min, respectively; Figure 6). A total of 4 red-tailed hawks, 2 northern harriers, 1 Swainson's hawk, 1 sharp-shinned hawk, were observed

at point 4 (Table 3). Species observed at point 2 included 2 northern harriers, 2 American kestrels, and 1 Swainson's hawk (Table 3). The habitat at point 2 and point 4 is not considered unique to the Project Area or present geographic features that would concentrate raptors to those specific areas.

3.2 FLIGHT HEIGHT AND ENCOUNTER RATE

During fall avian use surveys, biologists collected behavioral data for 99.9 percent of all birds observed during point count surveys of which 95.7 percent were observed flying. The biologists collected flight height data for 99.9 percent and flight direction for 47.6 percent of observations. Of the individual non-raptors observed flying, 49.0 percent flew below the height of the anticipated RSA, 39.7 percent flew at the height of the anticipated RSA, and 11.3 percent flew above the height of the anticipated RSA (Table 4). Of the individual raptors observed flying, 45.2 percent flew below the height of the anticipated RSA, 38.7 percent flew at the height of the anticipated RSA, and 16.1 percent flew above the height of the anticipated RSA (Table 4). Data on flight direction are located in Appendix 1. The majority of the observations were flying in a southerly or northerly direction.

The red-winged blackbird, unidentified blackbird, common grackle, snow goose, and horned lark had the highest encounter rates (7.16, 5.05, 1.19, 1.17, and 0.82 birds flying at RSA height/20 min, respectively; Table 5). All other species had encounter rates of 0.26 birds flying at RSA height/20 min or less.

3.3 LISTED SPECIES

No federally listed avian species were observed during fall avian surveys or as an incidental observation within the Project Area.

State-designated Species of Conservation Priority observed within the Project Area include grasshopper sparrow (level I), Swainson's hawk (level I), ferruginous hawk (priority I), northern harrier (level II), and sharp-tailed grouse (level II). Grasshopper sparrows (total of 11 individuals) were observed during point count surveys at all points except point 7. Swainson's hawks were observed at point 2 and point 4 (one individual at each) and as incidental observations. Northern harriers were observed at all points (total of 10 individuals) and as incidental observations. Two individual sharp-tailed grouse were observed at point 3. One ferruginous hawk was observed as an incidental observation.

3.6 INCIDENTAL OBSERVATIONS

Biologists documented 13 species as incidental observations (Table 6). Biologists documented three incidental species—ferruginous hawk, great horned owl, and gray partridge—that were not detected during fall point count surveys. Biologists observed several raptor species both as incidentals and during the point count surveys including the American kestrel, Swainson's hawk, northern harrier, and broad-winged hawk.

4.0 DISCUSSION

4.1 NON-RAPTOR USE AND ENCOUNTER RATE

The bird community of the Project Area was comprised primarily of grassland birds and shrubland birds that are commonly found in agricultural areas of the north central United States. The large flocks of snow geese observed in flight over the Project Area coincide with the annual fall migration in North Dakota for the species (USFWS 2011c). Red-winged blackbirds were the most commonly identified species within the Project Area. Most red-winged blackbirds were observed at point 4 which had a large field of planted sunflowers. Sunflowers are a favorite food source for the red-winged blackbird during the fall and winter seasons (Yasukawa and Searcy 1995). In North Dakota, red-winged blackbirds show a slight decreasing population trend (Sauer et al. 2011) but have a large population (11 million; Blancher et al. 2007). Red-winged blackbirds also had a high encounter rate and have been recorded as fatalities at other wind energy facilities but in low numbers (Kerlinger et al. 2006, Johnson and Erickson 2011). Given the low number of fatalities recorded at other wind farms, any fatalities should they occur in the Project Area are likely to be low and not expected to have impacts on the North Dakota population. Like the red-winged blackbird, common grackle (another blackbird) was a commonly identified species in the Project Area and had a high encounter rate. Mortality has been documented at other wind energy facilities for common grackle (Jain et al. 2007, and Johnson et al. 2002). In North Dakota, common grackle show an increasing population trend (Sauer et al. 2011) and have a large population (4.5 million; Blancher et al. 2007). Given the fatalities observed at other wind farms, any fatalities, should they occur at the Project, are not expected to have impacts on the North Dakota population. Horned larks were the second most common observed species in the Project Area and were observed at all survey points. Horned larks show a slight decreasing population trend (Sauer et al. 2011) but have a large population in North Dakota (6.8 million; Blancher et al. 2007). Horned larks had a high encounter rate and have been observed as fatalities at other wind energy projects (Derby et al. 2007, Stantec 2010, Johnson and Erickson 2011). As a result, fatalities of horned larks may occur in the Project Area but not expected to have impacts on the North Dakota population.

Snow geese were the third most common birds observed in the Project Area. During the fall, snow geese are commonly observed migrating through North Dakota on their way to wintering grounds in the southern part of the United States (USFWS 2011c). Results of the fall survey at the Project Area show that snow geese were observed in three large flocks (of up to several hundred birds each) flying in a southerly direction. None of the snow geese were observed on the ground or landing in the immediate vicinity of the Project Area. However, the snow geese did have a high encounter rate. The Mid-continental light geese (which include the snow geese) breeding population trend in Canada (which migrate through North Dakota) were estimated at over 3 million with a 19 percent increase from 2010 (USFWS 2011c). Overall, the mid-continental light geese population has shown a population increase dating back to 1970 (USFWS 2011c). Despite the abundance of snow geese, only one fatality has been documented at one wind energy facility (Anderson et al. 2005). Given the low amount of fatalities recorded at other wind energy facilities, it is anticipated that any fatalities of snow geese from the Project would be low. Additionally, North Dakota has a hunting season for snow geese in the fall from September 24 to December 30 (NDGF 2010).

4.2 RAPTOR USE AND ENCOUNTER RATE

High raptor use has been associated with high raptor mortality at wind farms (Erickson 2007). Conversely, raptor mortality appears to be low when raptor use is low, as defined by Erickson (2007) as less than 1.0 birds/20 min, which is the case for raptor use in the Project Area.

Red-tailed hawks and northern harriers were the most commonly observed raptor species during avian surveys. However, red-tailed hawks had a low encounter rate (0.08 birds flying at RSA height/20 min) and northern harriers had no encounter rate. As a result, any potential for turbine-related impacts to these species is anticipated to be low.

Swainson's hawks, American kestrels, sharp-shinned hawks, and broad winged hawks were also observed in the Project Area but with low encounter rates of 0.01 birds flying at RSA height/20 min (Table 5), thereby minimizing the potential for negative turbine-related impacts to these species. Mean use by raptors was highest at point count locations 2 and 4. However, the habitats at these points are not unique on the landscape regarding raptor species and these data should not be used to guide turbine siting discussions.

4.3 LISTED AND SENSITIVE SPECIES

No federally listed avian species were observed during fall avian surveys or as an incidental observation within the Project Area.

State-designated Species of Conservation Priority observed in the Project Area include grasshopper sparrow (priority I), Swainson's hawk (priority I), northern harrier (priority II), and sharp-tailed grouse (priority II). The designation of Species of Conservation Priority describes a species identified as in decline at the national, regional or state level, or a species whose population status is not well known, but thought to be in decline in North Dakota (Hagen et al. 2005). Species of Conservation Priority receive special attention from state agencies, but do not require take permits or have other regulatory implications regardless of status (Level I or II), but all of these species are protected under the Migratory Bird Treaty Act except sharp-tailed grouse. The sharp-tailed grouse is managed as a game-bird in North Dakota (NDGF 2010). All species listed above had no encounter rates within the anticipated RSA, primarily because of their low occurrence within the Project Area.

4.4 CONCLUSION

Fall non-raptor use in the Project Area was relatively high, primarily due to high use by red-winged blackbirds (and unidentified blackbirds which may include red-winged blackbird, common grackle, and brown-headed cowbird), common grackle, snow geese, and horned larks. These species have high encounter rates but also have large regional breeding populations in North Dakota (red-winged blackbirds, common grackle, and horned larks) and northern Canada (snow geese). If fatalities do occur they are unlikely to have population-level consequences. Nocturnal migrants (e.g., some songbirds) may pass through the Project Area and would not be detected by the survey methods used in this study if the birds did not stop over within the Project Area. However, mortality of nocturnal migrants from the Project is not expected to have population-level implications because less than 0.01 percent of nocturnal migrants that fly through wind farms are killed (Erickson 2007).

The level of raptor use in the Project Area suggests that raptor mortality is anticipated to be low, especially based on the results by Erickson (2007). Red-tailed hawks and northern harriers were the most common raptors observed at the Project but in low numbers. Red-tailed hawk had a low encounter rate and northern harrier had no encounter rate within the anticipated RSA. Any potential for turbine-related impacts to these species is anticipated to be low.

5.0 REFERENCES

- Anderson, R., J. Tom, N. Neumann, W. P. Erickson, M. D. Strickland, M. Bourasse, K. J. Bay, and K.J. Sernka. 2005. Avian monitoring and risk assessment at the San Gorgonio Wind Resource Area. Technical report prepared by State Energy Resources Conservation and Development Commission and Western EcoSystems Technology, Inc. for National Renewable Energy Laboratory. Golden, CO.
- Arnett, E.B., D.B. Inkley, D.H. Johnson, R.P. Larkin, S. Manes, A.M. Manville, J.R. Mason, M.L. Morrison, M.D. Strickland, and R. Thresher. 2007. Impacts of wind energy facilities on wildlife and wildlife habitat. Wildlife Society Technical Review 07-2. The Wildlife Society, Bethesda, MA.
- Ballam, J.M. 1984. The use of soaring by the red-tailed hawk (*Buteo jamaicensis*). *Auk* 3:519-524.
- Beveridge, L. J. 2005. The Migratory Bird Treaty Act and wind development. *North American Wind Power* September: 36-38.
- Blancher, P. J., K. V. Rosenberg, A. O. Panjabi, B. Altman, J. Bart, C. J. Beardmore, G. S. Butcher, D. Demarest, R. Dettmers, E. H. Dunn, W. Easton, W. C. Hunter, E. E. Inigo-Elias, D. N. Pashley, C. J. Ralph, T. D. Rich, C. M. Rustay, J. M. Ruth, and T. C. Will. 2007. Guide to the Partners in Flight Population Estimates Database. Version: North American Landbird Conservation Plan 2004. Partners in Flight Technical Series No 5. <http://www.partnersinflight.org/>. Accessed November 2011.
- Bryce, S.A., J.M. Omernik, D.A. Pater, M. Ulmer, J. Schaar, J. Freeouf, R. Johnson, P. Kuck, and S.H. Azevedo. 1996. Ecoregions of North Dakota and South Dakota, (color poster with map, descriptive text, summary tables, and photographs): Reston, Virginia, U.S. Geological Survey (map scale 1:1,500,000).
- Derby, C., A. Dahl, W. Erickson, K. Bay, and J. Hoban. 2007. Post-Construction Monitoring Report for Avian and Bat Mortality at the NPPD Ainsworth Wind Farm. Prepared by WEST, Inc. for Nebraska Public Power District.
- Drewitt, A.L., and R.H.W. Langston. 2006. Assessing the impacts of wind farms on birds. *Ibis* 148:29-42.
- Erickson, W.P. 2007. Summary of methods and results for prediction and estimation of impacts and risk. Presented at NWCC Probability of Impact Workshop, 13 November 2007, Golden, CO.
- Erickson, W.P., G.D. Johnson, and D.P. Young, Jr. 2005. A summary and comparison of bird mortality from anthropogenic causes with an emphasis on collisions. USDA Forest Service Gen. Tech. Rep. PSW-GTR-191.
- Erickson, W.P., J. Jeffrey, K. Kronner, and K. Bay. 2004. Stateline Wind Project Wildlife Monitoring Final Report, July 2001—December 2003. Technical report prepared for FPL Energy, the Oregon Energy Facility Siting Council, and the Stateline Technical Advisory Committee.
- Erickson, W.P., G. Johnson, D. Young, D. Strickland, R. Good, M. Bourassa, K. Bay, and K. Sernka. 2002. Synthesis and comparison of baseline avian and bat use, raptor nesting and

- mortality information from proposed and existing wind developments. Technical report prepared by WEST, Inc., for Bonneville Power Administration, Portland, OR.
- Erickson, W.P., G.D. Johnson, M.D. Strickland, D.P. Young Jr., K.J. Sernka, and R E. Good. 2001. Avian collisions with wind turbines: a summary of existing studies and comparisons to other sources of avian collision mortality in the United States. National Wind Coordinating Committee, Washington, DC. Accessed at: http://www.nationalwind.org/assets/archive/Avian_Collisions_with_Wind_Turbines_-_A_Summary_of_Existing_Studies_and_Comparisons_to_Other_Sources_of_Avian_Collision_Mortality_in_the_United_States__2001_.pdf.
- Faanes, C.A., and R.E. Stewart. 1982. Revised checklist of North Dakota birds. Version 16JUL97. *The Prairie Naturalist* 14(3):81-92. Jamestown, ND: Northern Prairie Wildlife Research Center Online. Available online at: <http://www.npwrc.usgs.gov/resource/birds/chnbird/chnbird.htm>. Accessed November 2011.
- Ferrer, M., M. de Lucas, G.F.E. Janss, E. Casado, A.R. Muñoz, M.J. Bechard, and C.P. Calabuig. 2011. Weak relationship between risk assessment studies and recorded mortality in wind farms. *Journal of Applied Ecology*. Published online September 2011. DOI 10.1111/j.1365-2664.2011.02054.x
- Hagen, S.K., P.T. Isakson, and S.R. Dyke. 2005. North Dakota Comprehensive Wildlife Conservation Strategy. North Dakota Game and Fish Department. Bismarck, ND.
- Jain, A., P. Kerlinger, R. Curry, and L. Slobodnik. 2007. Annual report for the Maple Ridge wind power project post-construction bird and bat fatality study—2006. Prepared by Curry and Kerlinger, LLC for PPM Energy, Horizon Energy, and Technical Advisory Committee for the Maple Ridge Project.
- Johnson, G.D. and W.P. Erickson. 2011. Avian, Bat and Habitat Cumulative Impacts Associated with Wind Energy Development in the Columbia Plateau Ecoregion of Eastern Washington and Oregon. Prepared for Klickitat County Planning Development.
- Johnson, G.D., W.P. Erickson, M.D. Strickland, M.F. Shepherd, D.A. Shepherd, and S.A. Sarappo. 2002. Collision mortality of local and migrant birds at a large-scale wind power development on Buffalo Ridge, Minnesota. *Wildlife Society Bulletin* 30:879-887.
- Kerlinger, P., R. Curry, L. Culp, A. Jain, C. Wilkerson, B. Fischer, and A. Hasch. 2006. Post construction avian and bat fatality monitoring study for the High Winds wind power project. Solano County, California: Two Year Report. Prepared for High Winds, LLC, FPL Energy.
- Kerns, J., and P. Kerlinger. 2004. A study of bird and bat collision fatalities at the Mountaineer Wind Energy Center, Tucker County, West Virginia: Annual report for 2003. Technical report prepared by Curry and Kerlinger, LLC for FPL Energy and Mountaineer Wind Energy Center Technical Review Committee.
- Leddy, K.L., K.F. Higgins, and D.E. Naugle. 1999. Effects of wind turbines on upland nesting birds in CRP grasslands. *Wilson Bulletin* 111:100-104.
- National Wind Coordinating Collaborative (NWCC). 2010. Wind Turbine Interactions with Birds and Bats: A Summary of Research Results and Priority Questions. Online at: <http://www.nationalwind.org/publications/bbfactsheet.aspx>. Accessed November 2011.

- North Dakota Game and Fish Department (NDGF). 2010. The 2011 Waterfowl Hunting Guide (Regulations). Available online at: <http://gf.nd.gov/regulations/waterfowl/index.html#light>. Accessed November 2011.
- Orloff, S., and A. Flannery. 1992. Wind turbine effects on avian activity, habitat use, and mortality in Altamont Pass and Solano County Wind Resource Areas, 1989-1991. Final report prepared by Biosystems Analysis, Inc. for Alameda, Contra Costa, and Solano Counties and the California Energy Commission.
- Pickwell, B. 1931. The prairie horned lark. *St. Louis Academy of Sciences Transactions* 27:1-153.
- Sauer, J.R., J.E. Hines, J.E. Fallon, K.L. Pardieck, D.J. Ziolkowski, Jr., and W.A. Link. 2011. The North American Breeding Bird Survey, Results and Analysis 1966 - 2009. Version 3.23.2011 USGS Patuxent Wildlife Research Center, Laurel, MD.
- Shaffer, J.A. and D.H. Johnson. 2008. Effects of wind developments on grassland birds in the northern Great Plains. NWCC Wind Wildlife Conference [Presentation], October 2008. Milwaukee, WI.
- Stantec Consulting Ltd. 2010. Wolfe Island Ecopower Center Post-Construction Follow-Up Plan Bird and Bat Resources: Monitoring Report No. 2 July-December 2009. Prepared for TransAlta Corporation's wholly owned subsidiary Canadian Renewable Energy Corporation.
- Strickland, M.D., E.B. Arnett, W.P. Erickson, D.H. Johnson, G.D. Johnson, M.L., Morrison, J.A. Shaffer, and W. Warren-Hicks. 2011. Comprehensive Guide to Studying Wind Energy/Wildlife Interactions. Prepared for the National Wind Coordinating Collaborative, Washington, D.C., USA. Available online at: http://www.nationalwind.org/assets/publications/Comprehensive_Guide_to_Studying_Wind_Energy_Wildlife_Interactions_2011_Updated.pdf. Accessed November 2011.
- Strickland, D., and M.L. Morrison. 2008. A summary of avian/wind facility interactions in the U.S. Federal Guidelines Committee for Wind Siting Guidelines, February 26, 2008, Washington, DC.
- Thompson, W.L. 2002. Towards reliable bird surveys: accounting for individuals present but not detected. *Auk* 119:18-25.
- USFWS (U.S. Fish and Wildlife Service). 2011a. Flyways. Retrieved from: <http://www.flyways.us/>. Accessed: November 2011.
- USFWS. 2011b. U.S. Fish and Wildlife Service Draft Land-Based Wind Energy Guidelines. Retrieved from: http://www.fws.gov/windenergy/docs/Wind_Energy_Guidelines_2_15_2011FINAL.pdf. Accessed November 2011.
- USFWS. 2011c. Waterfowl population status, 2011. U.S. Department of the Interior. Washington, D.C. USA.
- Yasukawa, K. and W.A. Searcy. 1995. Red-winged Blackbird (*Agelaius phoeniceus*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu.bnaproxy.birds.cornell.edu/bna/species/184.doi:10.2173/bna.184>. Accessed November 2011.

| | |
|------------------------|---|
| Jim Kowalsky | Nov. 18, 2011, revised June 8, 2012 |
| Report Author | Date |
| Sean Sparks | Nov. 21, 2011 |
| Peer Review #1 | Date |
| Lynn Sharpe | Nov. 22, 2011 |
| Peer Review #2 | Date |
| Tracey Dubuque | Nov. 28, 2011 |
| Project Manager | Date |
| Irina Gumennik | Nov. 25, 2011 |
| Deputy Project Manager | Date |
| Robert Friedel | Nov. 17, 2011 |
| GIS | Date |

FIGURES

Figure 2


Point count location map
(Fall 2011)



Wilton IV
Wind Energy Center

Burleigh County, ND

-  Avian survey point
-  Avian survey point
800-m radius
- PC# Point count number
-  Wind Energy Center
(11-15-2011)
-  Proposed turbine
(8-11-2011)
-  Local road



N 1:55,000
NAD 1983 UTM 14
Last modified: 11-16-2011



Miles

P:\GIS_PROJECTS\NextEra\Wilton_IV\MapDocs\Avian\2011\Fall\Report_Maps\NEER_Whitton_AvianSurvey_Figure2_PointCountLocation_17111_111611 - Last Accessed By: RF on 11/17/2011 - Map Scale: ANSIB (17" x 11")

Figure 3. Non-raptor mean use by survey date in Fall 2011 at the Wilton IV Wind Energy Center.




Figure 4

Non-raptor mean use by point count location (Fall 2011)



Wilton IV
Wind Eergy Center

Burleigh County, ND

Non-raptors per 20 minutes

- 0.01 - 25.00
- 25.01 - 50.00
- 50.01 - 75.00
- 75.01 - 100.00


Mean use value

PC# Point count number

Wind Energy Center (11-15-2011)

+ Proposed turbine (8-11-2011)

Local road



N
1:55,000
NAD 1983 UTM 14
Last modified: 11-16-2011




Figure 5. Raptor mean use by survey date in Fall 2011 at the Wilton IV Wind Energy Center.




Figure 6

Raptor mean use by point count location (Fall 2011)



Wilton IV
Wind Energy Center

Burleigh County, ND

Raptors per 20 minutes

- 0.01 - 0.21
- 0.22 - 0.42
- 0.43 - 0.62

Mean use value
PC# Point count number


Wind Energy Center (11-15-2011)

Proposed turbine (8-11-2011)

Local road

N 1:55,000
NAD 1983 UTM 14
Last modified: 11-16-2011

Miles



TABLES

Table 1. Fall 2011 point count survey dates at the Wilton IV Wind Energy Center.

| Survey number | Date(s) |
|----------------------|----------------|
| 1 | 8/9 |
| 2 | 8/17 |
| 3 | 8/24 |
| 4 | 9/2 |
| 5 | 9/9 |
| 6 | 9/16 |
| 7 | 9/23 |
| 8 | 9/30 |
| 9 | 10/7 |
| 10 | 10/15 |
| 11 | 10/20 |
| 12 | 10/28 |
| 13 | 11/3 |
| 14 | 11/11 |

Table 2. Avian species, by species grouping, observed during Fall 2011 point count surveys at the Wilton IV Wind Energy Center.

| Species Grouping | Overall Rank ¹ | Number of Birds | Number of Observations | Mean Use # birds per 20 min. (90% confidence interval) | Frequency % of surveys detected | Percent Composition | |
|------------------------|---------------------------|-----------------|------------------------|--|------------------------------------|---------------------|--------------|
| | | | | | | Group | Overall |
| Songbirds | | | | | | | |
| unidentified blackbird | 1 | 995 | 12 | 10.15 (2.24-18.06) | 10.2 | 30.0% | 24.4% |
| red-winged blackbird | 2 | 830 | 18 | 8.47 (0.00-18.55) | 14.3 | 25.1% | 20.3% |
| horned lark | 3 | 807 | 62 | 8.23 (4.73-11.73) | 40.8 | 24.4% | 19.8% |
| common grackle | 5 | 269 | 21 | 2.74 (0.60-4.88) | 14.3 | 8.1% | 6.6% |
| barn swallow | 6 | 135 | 30 | 1.38 (0.56-2.20) | 16.3 | 4.1% | 3.3% |
| American goldfinch | 7 | 90 | 39 | 0.92 (0.13-1.71) | 18.4 | 2.7% | 2.2% |
| American crow | 9 | 42 | 10 | 0.43 (0.12-0.74) | 9.2 | 1.3% | 1.0% |
| cedar waxwing | 10 | 35 | 1 | 0.36 (0.00-0.95) | 1.0 | 1.1% | 0.9% |
| western meadowlark | 13 | 23 | 12 | 0.23 (0.06-0.40) | 11.2 | 0.7% | 0.6% |
| European starling | 17 | 16 | 3 | 0.16 (0.00-0.35) | 2.0 | 0.5% | 0.4% |
| American robin | 18 | 15 | 5 | 0.15 (0.00-0.33) | 4.1 | 0.5% | 0.4% |
| house sparrow | 19 | 13 | 4 | 0.13 (0.00-0.27) | 3.1 | 0.4% | 0.3% |
| grasshopper sparrow | 21 | 11 | 11 | 0.11 (0.05-0.17) | 9.2 | 0.3% | 0.3% |
| vesper sparrow | 24 | 9 | 5 | 0.09 (0.01-0.17) | 5.1 | 0.3% | 0.2% |
| western kingbird | 25 | 6 | 6 | 0.06 (0.02-0.10) | 6.1 | 0.2% | 0.1% |
| tree swallow | 26 | 4 | 2 | 0.04 (0.00-0.11) | 1.0 | 0.1% | 0.1% |
| eastern kingbird | 26 | 4 | 3 | 0.04 (0.00-0.08) | 3.1 | 0.1% | 0.1% |
| American tree sparrow | 29 | 2 | 1 | 0.02 (0.00-0.05) | 1.0 | 0.1% | 0.0% |
| yellow warbler | 34 | 1 | 1 | 0.01 (0.00-0.03) | 1.0 | 0.0% | 0.0% |
| northern shrike | 34 | 1 | 1 | 0.01 (0.00-0.03) | 1.0 | 0.0% | 0.0% |
| eastern bluebird | 34 | 1 | 1 | 0.01 (0.00-0.03) | 1.0 | 0.0% | 0.0% |
| chipping sparrow | 34 | 1 | 1 | 0.01 (0.00-0.03) | 1.0 | 0.0% | 0.0% |
| brown thrasher | 34 | 1 | 1 | 0.01 (0.00-0.03) | 1.0 | 0.0% | 0.0% |
| blue jay | 34 | 1 | 1 | 0.01 (0.00-0.03) | 1.0 | 0.0% | 0.0% |
| Group Total | | 3312 | 251 | 33.80 (19.27-48.33) | 84.7 | | 81.1% |
| Waterfowl | | | | | | | |
| snow goose | 4 | 540 | 3 | 5.51 (0.09-10.93) | 3.1 | 93.3% | 13.2% |
| Canada goose | 12 | 28 | 3 | 0.29 (0.00-0.70) | 3.1 | 4.8% | 0.7% |
| mallard | 21 | 11 | 2 | 0.11 (0.00-0.24) | 2.0 | 1.9% | 0.3% |
| Group Total | | 579 | 8 | 5.91 (0.48-11.34) | 8.2 | | 14.2% |

Table 2. Avian species, by species grouping, observed during Fall 2011 point count surveys at the Wilton IV Wind Energy Center.

| Species Grouping | Overall Rank ¹ | Number of Birds | Number of Observations | Mean Use # birds per 20 min. (90% confidence interval) | Frequency % of surveys detected | Percent Composition | |
|----------------------|---------------------------|-----------------|------------------------|--|------------------------------------|---------------------|-------------|
| | | | | | | Group | Overall |
| Pigeons/Doves | | | | | | | |
| mourning dove | 8 | 67 | 33 | 0.68 (0.33-1.03) | 25.5 | 78.8% | 1.6% |
| rock pigeon | 15 | 18 | 5 | 0.18 (0.04-0.32) | 5.1 | 21.2% | 0.4% |
| Group Total | | 85 | 38 | 0.87 (0.51-1.23) | 29.6 | | 2.1% |
| Raptors | | | | | | | |
| red-tailed hawk | 16 | 17 | 16 | 0.17 (0.09-0.25) | 14.3 | 50.0% | 0.4% |
| northern harrier | 23 | 10 | 10 | 0.10 (0.05-0.15) | 10.2 | 29.4% | 0.2% |
| Swainson's hawk | 29 | 2 | 2 | 0.02 (0.00-0.04) | 2.0 | 5.9% | 0.0% |
| American kestrel | 29 | 2 | 2 | 0.02 (0.00-0.04) | 2.0 | 5.9% | 0.0% |
| unidentified buteo | 34 | 1 | 1 | 0.01 (0.00-0.03) | 1.0 | 2.9% | 0.0% |
| sharp-shinned hawk | 34 | 1 | 1 | 0.01 (0.00-0.03) | 1.0 | 2.9% | 0.0% |
| broad-winged hawk | 34 | 1 | 1 | 0.01 (0.00-0.03) | 1.0 | 2.9% | 0.0% |
| Group Total | | 34 | 33 | 0.35 (0.25-0.45) | 27.6 | | 0.8% |
| Gamebirds | | | | | | | |
| ring-necked pheasant | 11 | 30 | 27 | 0.31 (0.18-0.44) | 22.4 | 93.8% | 0.7% |
| sharp-tailed grouse | 29 | 2 | 1 | 0.02 (0.00-0.05) | 1.0 | 6.3% | 0.0% |
| Group Total | | 32 | 28 | 0.33 (0.20-0.46) | 23.5 | | 0.8% |
| Waterbirds | | | | | | | |
| American coot | 14 | 21 | 2 | 0.21 (0.00-0.55) | 2.0 | 80.8% | 0.5% |
| killdeer | 26 | 4 | 2 | 0.04 (0.00-0.09) | 2.0 | 15.4% | 0.1% |
| great blue heron | 34 | 1 | 1 | 0.01 (0.00-0.03) | 1.0 | 3.8% | 0.0% |
| Group Total | | 26 | 5 | 0.27 (0.00-0.61) | 5.1 | | 0.6% |
| Cranes/Rails | | | | | | | |
| sandhill crane | 20 | 12 | 1 | 0.12 (0.00-0.32) | 1.0 | 100.0% | 0.3% |
| Group Total | | 12 | 1 | 0.12 (0.00-0.32) | 1.0 | | 0.3% |
| Woodpeckers | | | | | | | |
| northern flicker | 29 | 2 | 2 | 0.02 (0.00-0.04) | 2.0 | 66.7% | 0.0% |
| hairy woodpecker | 34 | 1 | 1 | 0.01 (0.00-0.03) | 1.0 | 33.3% | 0.0% |
| Group Total | | 3 | 3 | 0.03 (0.00-0.06) | 3.1 | | 0.1% |
| Grand Total | | 4083 | 367 | 41.66 (26.10-57.22) | | | |

¹ A ranking of 1 indicates highest mean use

Table 3. Avian species observed by point during Fall 2011 point count surveys at the Wilton IV Wind Energy Center.

| Species | Number of Birds | Number of Obs. | Points | | | | | | |
|------------------------|-----------------|----------------|--------|-----|----|-----|----|-----|-----|
| | | | 1 | 2 | 3 | 4 | 5 | 6 | 7 |
| unidentified blackbird | 995 | 12 | 18 | 265 | 40 | 298 | 24 | 0 | 350 |
| red-winged blackbird | 830 | 18 | 6 | 111 | 38 | 633 | 34 | 0 | 8 |
| horned lark | 807 | 62 | 11 | 232 | 82 | 73 | 78 | 44 | 287 |
| snow goose | 540 | 3 | 115 | 0 | 0 | 0 | 0 | 175 | 250 |
| common grackle | 269 | 21 | 18 | 118 | 36 | 40 | 4 | 45 | 8 |
| barn swallow | 135 | 30 | 1 | 103 | 1 | 21 | 0 | 4 | 5 |
| American goldfinch | 90 | 39 | 8 | 60 | 6 | 11 | 2 | 1 | 2 |
| mourning dove | 67 | 33 | 10 | 12 | 3 | 11 | 3 | 5 | 23 |
| American crow | 42 | 10 | 12 | 6 | 7 | 16 | 0 | 1 | 0 |
| cedar waxwing | 35 | 1 | 0 | 35 | 0 | 0 | 0 | 0 | 0 |
| ring-necked pheasant | 30 | 27 | 2 | 4 | 4 | 3 | 3 | 9 | 5 |
| Canada goose | 28 | 3 | 0 | 0 | 0 | 1 | 27 | 0 | 0 |
| western meadowlark | 23 | 12 | 1 | 9 | 3 | 0 | 1 | 1 | 8 |
| American coot | 21 | 2 | 0 | 0 | 0 | 1 | 0 | 0 | 20 |
| rock pigeon | 18 | 5 | 0 | 0 | 0 | 0 | 3 | 15 | 0 |
| red-tailed hawk | 17 | 16 | 3 | 0 | 4 | 4 | 2 | 2 | 2 |
| European starling | 16 | 3 | 0 | 7 | 0 | 0 | 9 | 0 | 0 |
| American robin | 15 | 5 | 0 | 15 | 0 | 0 | 0 | 0 | 0 |
| house sparrow | 13 | 4 | 0 | 13 | 0 | 0 | 0 | 0 | 0 |
| sandhill crane | 12 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 12 |
| mallard | 11 | 2 | 0 | 0 | 0 | 11 | 0 | 0 | 0 |
| grasshopper sparrow | 11 | 11 | 1 | 1 | 2 | 2 | 2 | 3 | 0 |
| northern harrier | 10 | 10 | 1 | 2 | 1 | 2 | 1 | 1 | 2 |
| vesper sparrow | 9 | 5 | 0 | 0 | 0 | 0 | 3 | 1 | 5 |
| western kingbird | 6 | 6 | 1 | 1 | 1 | 0 | 1 | 2 | 0 |
| tree swallow | 4 | 2 | 0 | 0 | 4 | 0 | 0 | 0 | 0 |
| killdeer | 4 | 2 | 0 | 2 | 2 | 0 | 0 | 0 | 0 |
| eastern kingbird | 4 | 3 | 0 | 3 | 0 | 0 | 0 | 1 | 0 |
| Swainson's hawk | 2 | 2 | 0 | 1 | 0 | 1 | 0 | 0 | 0 |
| sharp-tailed grouse | 2 | 1 | 0 | 0 | 2 | 0 | 0 | 0 | 0 |
| northern flicker | 2 | 2 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| American tree sparrow | 2 | 1 | 0 | 2 | 0 | 0 | 0 | 0 | 0 |
| American kestrel | 2 | 2 | 0 | 2 | 0 | 0 | 0 | 0 | 0 |
| yellow warbler | 1 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 |
| unidentified buteo | 1 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 |
| sharp-shinned hawk | 1 | 1 | 0 | 0 | 0 | 1 | 0 | 0 | 0 |
| northern shrike | 1 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 |
| hairy woodpecker | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 0 | 0 |
| great blue heron | 1 | 1 | 0 | 0 | 0 | 1 | 0 | 0 | 0 |
| eastern bluebird | 1 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 |

Table 3. Avian species observed by point during Fall 2011 point count surveys at the Wilton IV Wind Energy Center.

| Species | Number of Birds | Number of Obs. | Points | | | | | | |
|--------------------|-----------------|----------------|------------|-------------|------------|-------------|------------|------------|------------|
| | | | 1 | 2 | 3 | 4 | 5 | 6 | 7 |
| chipping sparrow | 1 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 |
| broad-winged hawk | 1 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 |
| brown thrasher | 1 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 |
| blue jay | 1 | 1 | 0 | 0 | 0 | 0 | 1 | 0 | 0 |
| Grand Total | 4083 | 367 | 210 | 1010 | 237 | 1130 | 199 | 310 | 987 |

Table 4. Summary of avian flight heights¹ in relation to the turbine rotor swept area (RSA)² during Fall 2011 point count surveys at the Wilton IV Wind Energy Center.

| | Birds | |
|--------------------------|---------------|-------------------|
| | Number | Percentage |
| Non-raptors | | |
| Above RSA height (>130m) | 437 | 11.3% |
| At RSA height (30m–130m) | 1537 | 39.7% |
| Below RSA height (<30m) | 1896 | 49.0% |
| Raptors | | |
| Above RSA height (>130m) | 5 | 16.1% |
| At RSA height (30m–130m) | 12 | 38.7% |
| Below RSA height (<30m) | 14 | 45.2% |

¹ Includes only flying birds with flight height data

² These values assume a rotor diameter of 100 (m) and a hub height of 80 (m)

Table 5. Avian flight height characteristics in relation to the turbine rotor swept area (RSA)¹ during Fall 2011 point count surveys at the Wilton IV Wind Energy Center.

| Species | Encounter Rate | Mean Use # birds/ 20 min. (90% confidence interval) | Percent Flying | Percent Above RSA Height | Percent At RSA Height | Percent Below RSA Height |
|------------------------|-----------------------|--|-----------------------|---------------------------------|------------------------------|---------------------------------|
| red-winged blackbird | 7.16 | 8.47 (0.00 - 18.55) | 100.0 | 0.0 | 84.6 | 15.4 |
| unidentified blackbird | 5.05 | 10.15 (2.24 - 18.06) | 95.4 | 0.0 | 52.2 | 47.8 |
| common grackle | 1.19 | 2.74 (0.60 - 4.88) | 99.6 | 0.0 | 43.7 | 56.3 |
| snow goose | 1.17 | 5.51 (0.09 - 10.93) | 100.0 | 78.7 | 21.3 | 0.0 |
| horned lark | 0.82 | 8.23 (4.73 - 11.73) | 99.0 | 0.0 | 10.0 | 90.0 |
| Canada goose | 0.26 | 0.29 (0.00 - 0.70) | 100.0 | 0.0 | 89.3 | 10.7 |
| red-tailed hawk | 0.08 | 0.17 (0.09 - 0.25) | 88.2 | 33.3 | 53.3 | 13.3 |
| killdeer | 0.02 | 0.04 (0.00 - 0.09) | 100.0 | 0.0 | 50.0 | 50.0 |
| American crow | 0.01 | 0.43 (0.12 - 0.74) | 95.2 | 0.0 | 2.5 | 97.5 |
| unidentified buteo | 0.01 | 0.01 (0.00 - 0.03) | 100.0 | 0.0 | 100.0 | 0.0 |
| Swainson's hawk | 0.01 | 0.02 (0.00 - 0.04) | 100.0 | 0.0 | 50.0 | 50.0 |
| sharp-shinned hawk | 0.01 | 0.01 (0.00 - 0.03) | 100.0 | 0.0 | 100.0 | 0.0 |
| broad-winged hawk | 0.01 | 0.01 (0.00 - 0.03) | 100.0 | 0.0 | 100.0 | 0.0 |
| yellow warbler | 0.00 | 0.01 (0.00 - 0.03) | 100.0 | 0.0 | 0.0 | 100.0 |
| western meadowlark | 0.00 | 0.23 (0.06 - 0.40) | 39.1 | 0.0 | 0.0 | 100.0 |
| western kingbird | 0.00 | 0.06 (0.02 - 0.10) | 66.7 | 0.0 | 0.0 | 100.0 |
| vesper sparrow | 0.00 | 0.09 (0.01 - 0.17) | 33.3 | 0.0 | 0.0 | 100.0 |
| tree swallow | 0.00 | 0.04 (0.00 - 0.11) | 100.0 | 0.0 | 0.0 | 100.0 |
| sharp-tailed grouse | 0.00 | 0.02 (0.00 - 0.05) | 100.0 | 0.0 | 0.0 | 100.0 |
| sandhill crane | 0.00 | 0.12 (0.00 - 0.32) | 100.0 | 100.0 | 0.0 | 0.0 |
| rock pigeon | 0.00 | 0.18 (0.04 - 0.32) | 100.0 | 0.0 | 0.0 | 100.0 |
| ring-necked pheasant | 0.00 | 0.31 (0.18 - 0.44) | 3.3 | 0.0 | 0.0 | 100.0 |
| northern shrike | 0.00 | 0.01 (0.00 - 0.03) | 0.0 | 0.0 | 0.0 | 0.0 |
| northern harrier | 0.00 | 0.10 (0.05 - 0.15) | 100.0 | 0.0 | 0.0 | 100.0 |
| northern flicker | 0.00 | 0.02 (0.00 - 0.04) | 50.0 | 0.0 | 0.0 | 100.0 |
| mourning dove | 0.00 | 0.68 (0.33 - 1.03) | 59.7 | 0.0 | 0.0 | 100.0 |

Table 5. Avian flight height characteristics in relation to the turbine rotor swept area (RSA)¹ during Fall 2011 point count surveys at the Wilton IV Wind Energy Center.

| Species | Encounter Rate | Mean Use # birds/ 20 min. (90% confidence interval) | Percent Flying | Percent Above RSA Height | Percent At RSA Height | Percent Below RSA Height |
|-----------------------|-----------------------|--|-----------------------|---------------------------------|------------------------------|---------------------------------|
| mallard | 0.00 | 0.11 (0.00 - 0.24) | 100.0 | 0.0 | 0.0 | 100.0 |
| house sparrow | 0.00 | 0.13 (0.00 - 0.27) | 76.9 | 0.0 | 0.0 | 100.0 |
| hairy woodpecker | 0.00 | 0.01 (0.00 - 0.03) | 0.0 | 0.0 | 0.0 | 0.0 |
| grasshopper sparrow | 0.00 | 0.11 (0.05 - 0.17) | 0.0 | 0.0 | 0.0 | 0.0 |
| great blue heron | 0.00 | 0.01 (0.00 - 0.03) | 100.0 | 0.0 | 0.0 | 100.0 |
| European starling | 0.00 | 0.16 (0.00 - 0.35) | 100.0 | 0.0 | 0.0 | 100.0 |
| eastern kingbird | 0.00 | 0.04 (0.00 - 0.08) | 100.0 | 0.0 | 0.0 | 100.0 |
| eastern bluebird | 0.00 | 0.01 (0.00 - 0.03) | 0.0 | 0.0 | 0.0 | 0.0 |
| chipping sparrow | 0.00 | 0.01 (0.00 - 0.03) | 0.0 | 0.0 | 0.0 | 0.0 |
| cedar waxwing | 0.00 | 0.36 (0.00 - 0.95) | 100.0 | 0.0 | 0.0 | 100.0 |
| brown thrasher | 0.00 | 0.01 (0.00 - 0.03) | 100.0 | 0.0 | 0.0 | 100.0 |
| blue jay | 0.00 | 0.01 (0.00 - 0.03) | 0.0 | 0.0 | 0.0 | 0.0 |
| barn swallow | 0.00 | 1.38 (0.56 - 2.20) | 100.0 | 0.0 | 0.0 | 100.0 |
| American tree sparrow | 0.00 | 0.02 (0.00 - 0.05) | 100.0 | 0.0 | 0.0 | 100.0 |
| American robin | 0.00 | 0.15 (0.00 - 0.33) | 100.0 | 0.0 | 0.0 | 100.0 |
| American kestrel | 0.00 | 0.02 (0.00 - 0.04) | 50.0 | 0.0 | 0.0 | 100.0 |
| American goldfinch | 0.00 | 0.92 (0.13 - 1.71) | 96.7 | 0.0 | 0.0 | 100.0 |
| American coot | 0.00 | 0.21 (0.00 - 0.55) | 0.0 | 0.0 | 0.0 | 0.0 |

¹These values assume a rotor diameter of 100 (m) and a hub height of 80 (m)

Table 6. Incidental observations of birds during Fall 2011 point count surveys at the Wilton IV Wind Energy Center.

| Species |
|--------------------|
| American crow |
| American kestrel |
| American robin |
| broad-winged hawk |
| European starling |
| ferruginous hawk |
| great horned owl |
| gray partridge |
| mallard |
| northern harrier |
| red-tailed hawk |
| Swainson's hawk |
| western meadowlark |

APPENDICES

Appendix 1. Flight directions of birds observed during Fall 2011 point count surveys at the Wilton IV Wind Energy Center.

| Species | Number of Birds ¹ | Number of Observations | Percentage of Flights | | | | | | | | |
|------------------------|------------------------------|------------------------|-----------------------|------------|------------|-------------|-------------|------------|------------|-------------|------------|
| | | | N | NE | E | SE | S | SW | W | NW | Variable |
| red-winged blackbird | 737 | 11 | 67.8 | 0.0 | 0.0 | 0.0 | 12.9 | 0.0 | 0.0 | 14.9 | 0.0 |
| snow goose | 540 | 3 | 0.0 | 0.0 | 0.0 | 67.6 | 32.4 | 0.0 | 0.0 | 0.0 | 0.0 |
| unidentified blackbird | 400 | 2 | 37.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 62.5 | 0.0 |
| American crow | 35 | 5 | 0.0 | 0.0 | 0.0 | 60.0 | 5.7 | 8.6 | 0.0 | 0.0 | 0.0 |
| Canada goose | 28 | 3 | 0.0 | 14.3 | 0.0 | 85.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| horned lark | 21 | 9 | 19.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 19.0 | 14.3 | 28.6 |
| red-tailed hawk | 15 | 14 | 20.0 | 20.0 | 0.0 | 26.7 | 13.3 | 6.7 | 13.3 | 0.0 | 0.0 |
| mourning dove | 13 | 5 | 0.0 | 0.0 | 0.0 | 0.0 | 7.7 | 0.0 | 0.0 | 0.0 | 0.0 |
| sandhill crane | 12 | 1 | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| rock pigeon | 11 | 3 | 0.0 | 0.0 | 0.0 | 72.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| barn swallow | 10 | 4 | 0.0 | 30.0 | 0.0 | 0.0 | 10.0 | 0.0 | 0.0 | 30.0 | 0.0 |
| American robin | 10 | 1 | 100.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| northern harrier | 9 | 9 | 22.2 | 0.0 | 11.1 | 11.1 | 44.4 | 0.0 | 11.1 | 0.0 | 0.0 |
| American goldfinch | 6 | 6 | 0.0 | 0.0 | 0.0 | 16.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| common grackle | 3 | 1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| western kingbird | 2 | 2 | 0.0 | 0.0 | 0.0 | 0.0 | 50.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Swainson's hawk | 2 | 2 | 0.0 | 0.0 | 0.0 | 0.0 | 50.0 | 0.0 | 0.0 | 50.0 | 0.0 |
| American tree sparrow | 2 | 1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 100.0 |
| unidentified buteo | 1 | 1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 | 0.0 |
| sharp-shinned hawk | 1 | 1 | 0.0 | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| eastern kingbird | 1 | 1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| broad-winged hawk | 1 | 1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 | 0.0 | 0.0 |
| Grand Total | 1860 | 86 | 36.0 | 0.5 | 0.1 | 23.4 | 15.2 | 0.3 | 0.4 | 19.7 | 0.4 |

¹ Includes only flying birds with flight directions